

# STIC Search Report

## Biotech-Chem Library

STIC Database Tracking Number: 169018

TO: Ilia Ouspenski  
Location: REM-3D74&3C70  
Art Unit: 1644  
Thursday, October 20, 2005

Case Serial Number: 10/625105

From: Deirdre Arnold  
Location: Biotech-Chem Library  
REM 1A64  
Phone: 571-272-2532

Deirdre.Arnold@uspto.gov

### Search Notes

**RUSH**

*Please feel free to contact me if you have any questions or would like to amend the search.*

Thank you for using STIC services.

Regards,  
Deirdre Arnold



4  
STIC-Biotech/ChemLib

169018

From: Chan, Christina  
Sent: Wednesday, October 19, 2005 2:51 PM  
To: Ouspenski, Ilia; STIC-Biotech/ChemLib  
Subject: RE: RUSH sequence search request for 10/625,105

Please rush. Thanks Chris

Chris Chan  
TC 1600 New Hire Training Coordinator and SPE 1644  
(571)-272-0841  
Remsen, 3E89

(RFE)

-----Original Message-----

From: Ouspenski, Ilia  
Sent: Wednesday, October 19, 2005 2:29 PM  
To: Chan, Christina  
Subject: RUSH sequence search request for 10/625,105

Christina, please approve a RUSH search:

STIC:

please provide an INTERFERENCE ONLY search for amino acid sequences SEQ ID NOS: 28, 30, 32, 34, 36, and 38 for 10/625,105.

Thnaks,

ILIA OUSPENSKI, Ph.D.  
Examiner  
Art Unit 1644  
Phone: 571-272-2920  
REM 3D74  
Mailstop 3c70

\*\*\*\*\*

Searcher: \_\_\_\_\_  
Searcher Phone: \_\_\_\_\_  
Date Searcher Picked up: \_\_\_\_\_  
Date completed: \_\_\_\_\_  
Searcher Prep Time: \_\_\_\_\_  
Online Time: \_\_\_\_\_

\*\*\*\*\*

Type of Search  
NA# \_\_\_\_\_ AA# \_\_\_\_\_  
S/L: \_\_\_\_\_ Oligomer: \_\_\_\_\_  
Encode/Transl: \_\_\_\_\_  
Structure #: \_\_\_\_\_ Text: \_\_\_\_\_  
Inventor: \_\_\_\_\_ Litigation: \_\_\_\_\_

\*\*\*\*\*

Vendors and cost where applicable  
STN: \_\_\_\_\_  
DIALOG: \_\_\_\_\_  
QUESTEL/ORBIT: \_\_\_\_\_  
LEXIS/NEXIS: \_\_\_\_\_  
SEQUENCE SYSTEM: \_\_\_\_\_  
WWW/Internet: \_\_\_\_\_  
Other (Specify): \_\_\_\_\_

Tezuka et al., Poster, 1994.

**OBJECTIVE:** Recently, adhesion molecules in immune system cells have been identified and their functions have been clarified, but when one considers the complex nature of the immune system and tissue specificities and so forth, it is presumed that unknown adhesion molecules exist. The JTT.1 antibody, which we obtained in the course of searching for novel adhesion molecules, has a function to aggregate rat thymoma cell line FTL435 cells via an unknown adhesion pathway. Therefore it was believed the antigen molecule recognized by the JTT.1 antibody was a signal transduction molecule that induces an unknown cell adhesion. This time we produced the antibody JTT.2 with adhesion blocking activity and cloned the JTT.1 antigen gene in order to further clarify the properties of this adhesion phenomenon. As a result, we report that it was clarified that the JTT.1 antigen not only functions as a signal transduction molecule, but is also an adhesion molecule.

**METHOD:** Antibody preparation: Made by immunizing BALB/c mice with rat thymoma cell line FTL435, and fusing lymph node cells with mouse myeloma cells. Immunoprecipitation: Performed by making biotinylated cells soluble and using antibody-binding beads. Precipitates were subjected to SDS-PAGE, then transferred to a film and detected using an ECL system. Genetic cloning: poly(A)<sup>+</sup> RNA was prepared from rat ConA blast, cDNA was synthesized with this as a template, and then it was incorporated into an expression vector and a cDNA library was constructed. The library was transiently expressed with COS cells, and screened by panning using the JTT.1 antibody. The cDNA sequences concentrated by panning were determined by the dideoxy method. Flow cytometry analysis: Cells were treated with the antibody, stained with FITC-conjugated anti-mouse Ig, and then analyzed by flow cytometry.

**RESULTS AND CONSIDERATIONS:** A mouse was immunized with FTL435 cells, monoclonal antibodies were prepared in accordance with an index for blocking aggregation of the FTL435 cells by JTT.1 antibody, and JTT.2 was obtained. The isotype of JTT.2 antibody was IgG1κ. The JTT.2 antigen was strongly expressed in FTL435 cells, thymocytes, and activated lymphocytes. The result of immunoprecipitation indicated that the molecular weight of the JTT.2 antigen molecule was 24 kDa and 28 kDa by reduction. This matched the case of the JTT.1 antigen's molecule. Based on this, it was believed the JTT.2 antibody recognized the JTT.1 antigen. Therefore we performed an experiment to bind rat thymocytes to the JTT.1 antigen immobilized on a plate. Thymocytes in an unstimulated state did not bind to the JTT.1 antigen, but bound specifically to the JTT.1 antigen on a plate when stimulated with the JTT.1 antibody. This adhesion phenomenon was blocked by adding the JTT.2 antibody. Based on these results, it was believed the JTT.1 antigen is an adhesion molecule.

Next, we performed genetic cloning of the JTT.1 antigen. A cDNA library was prepared from rat ConA blasts that express the JTT.1 antigen. The cDNA library was transiently expressed in COS cells, and panning using the JTT.1 antibody was repeated three times, thus cloning a 0.9 kbp gene. COS cells that transiently expressed this gene reacted strongly with the JTT.1 and JTT.2 antibodies. A sequence matching the N-terminal sequence of the purified JTT.1 antigen was present in the amino acid sequence predicted from the nucleotide sequence. Based on these results, it was believed that the gene we cloned was the JTT.1 antigen gene. In the predicted amino acid sequence there

were sugar-chain binding sites at two positions in the extracellular domain, and, in the intracellular domain, there was a PKC phosphorylation site at one position and CK2 phosphorylation sites at two positions. Furthermore, we performed homology searching for the entire length of the amino acid sequence, but no homologous molecule among previously known molecules was found. Therefore it is believed the JTT.1 antigen is a novel adhesion molecule.

#### SUMMARY

1. A JTT.2 antibody that blocks unknown cell aggregation induced by the JTT.1 antigen was prepared.
2. It was believed the JTT.2 antibody recognizes the JTT.1 antigen.
3. Rat thymocytes stimulated by the JTT.1 antigen bound to the JTT.1 antigen. This binding was specifically blocked by the JTT.2 antibody.
4. The JTT.1 antigen gene was obtained by expression cloning. Homology searching resulted in finding no highly similar molecule.

Based on these results, it is believed the JTT.1 antigen is a novel adhesion molecule.





FIG. 1  
JTT.2 antibody's FTL435 cellular aggregation blocking activity.  
FTL435 cells were cultured at 37 degrees C for one hour in the presence or absence of the antibody.

**FIG. 2**

Comparison of expression patterns of JTT.1 antigen and JTT.2 antigen in various cells. FTL435, thymocytes, splenic lymphocytes, and ConA blasts were stained with antibodies and analyzed with flow cytometry.





FIG. 3


Comparison of JTT.1 antigen and JTT.2 antigen according to immunoprecipitation.

Each antigen was immunoprecipitated using the soluble material of biotinylated FTL435 cells, subjected to SDS-PAGE, transferred to a film, and detected with an ECL system.



**FIG. 4**  
The JTT.2 antigen blocks binding of thymocytes to the JTT.1 antigen.  
Purified JTT.1 antigen was coated on a plate;  
rat thymocytes were cultured at 37 degrees C for one hour  
in the presence and absence of the antibody.





**FIG. 5**  
**JTT.1 antigen expression cloning.**

(A) Summary of cDNA cloning using panning method.  
(B) Analysis of plasmids recovered using COS cells: Restriction enzyme digestion of recovered plasmids, then 1% agarose gel electrophoresis, and analyze insert DNA.  
(C) Reactivity of JTT.1 antigen cDNA product with JTT.1 antibody: Use JTT.1 antibody to stain COS cells that transiently expressed JTT.1 antigen cDNA obtained by panning, and analyze with flow cytometry.

## 【目的】

最近、免疫系細胞における抗原分子の同定や機能解明が行われているが、免疫系の多様性や組織特異性などを考えると未知の抗原分子の存在が予想される。われわれが新たに抗原分子を探索する過程で得たJIT.1抗体は、ラット胸腺腫細胞株 FTL435 細胞を未知の抗原細胞を用いて感染させる作用を有する。このことから、JIT.1抗体が認識している抗原分子は、未知の細胞抗原を誘導するシグナル伝達分子と考えられた。今回、この抗原分子の性状をさらに明らかにするために、抗原阻害性を有する抗体JIT.2を作成するとともに、JIT.1抗原遺伝子をクローニングした。その結果、JIT.1抗原がシグナル伝達分子としての機能を有するだけでなく、抗原分子であることが明らかとなり、その報告する。

【方法】 抗体の作成：ラット胸腺リンパ節細胞をラウスニエロー細胞と沈降：デオキシ化した細胞を可溶化し、SDS-PAGE後膜に転写し、ECLシステム、ラット Con A blast より poly(A)<sup>+</sup>RNA を発現ヘクターに組み込み cDNA ライブラリーで一過性発現させ、JIT1 抗体を用いたハイニグにより選別された cDNA の遺伝子クローニングを行った。細胞を抗体で染色後、FACS 分析を行った。



## まとめ

1. JTT.1抗体により除去される未知の抗原と阻害するJTT.2抗体を単製した。  
と考えられた。
  2. JTT.2抗体は、JTT.1抗原を認識し、抗原に結合した。
  3. JTT.1抗体刺激したマウス脾臓細胞の培養上清は、JTT.2抗体により阻害された。
  4. JTT.1抗体を除去する処理のロケータは確立された。
- 以上の結果からJTT.1抗原は、新規の抗原であると考えられる。




図1. JTT.2抗体のFTL435細胞集阻害活性。  
FTL435細胞を抗体存在下あるいは非存在  
下で37℃、1h培養した。





図2. 各種細胞におけるJTT.1抗原とJTT.2抗原の発現パターンの比較。

FTL435、胸腺細胞、脾臓リンパ球、ConA blast を抗体で染色し、フローサイトメーターで解析した。








図 4. JTT.2抗体は胸腺細胞のJTT.1抗原への結合を阻害する。  
精製JTT.1抗原をプレートにコートし、抗体存在下あるいは非存在下でラット胸腺細胞を37℃、1h培養した。





図5. JTT.1抗原の発現クローニング。

- (A) パニング法によるcDNAクローニングの概略
- (B) COS細胞より回収されたプラスミドの解析：回収されたプラスミドを制限酵素消化後、1%アガロースゲル電気泳動を行い、インサートDNAを解析した。
- (C) JTT.1抗原cDNA産物とJTT.1抗体との反応性：パニング法により得られたJTT.1抗原cDNAを一過性発現させたCOS細胞をJTT.1抗体で染色し、フローサイトメーターで解析した。



## TRANSLATION from Japanese to English

## CERTIFICATE OF ACCURACY

This day personally appeared before me Gregor Hartmann, who after being duly sworn deposes and states:

that he is a translator of the Japanese and English languages, a professional provider of translations, accredited by the American Translators Association for Japanese to English translation;

that he is thoroughly familiar with these languages and has carefully made and verified the attached translation from the original document in the Japanese language, to wit:

Untitled article on research on monoclonal antibodies (JTT.1, JTT.2)

and that the attached translation is a true and correct English version of the original to the best of his knowledge and belief.

Gregor Hartmann  
Gregor Hartmann, Translator

Sworn to before me

this date: 12/27/02

Gene Bressler  
Notary Public

Gene Bressler  
Notary Public of New Jersey  
My Commission Expires 5/30/2007